

APPLICATION
FOR
UNITED STATES PATENT

**A DEVICE FOR DETECTING BACTERIAL CONTAMINATION
AND METHOD OF USE**

Inventor:

MITCHELL C. SANDERS

NUTTER, McCLENNEN & FISH, LLP
One International Place
Boston, MA 02110-2699
Telephone (617) 439-2000
Facsimile (617) 973-9748

EXPRESS MAIL NO.: EL835814223US

A DEVICE FOR DETECTING BACTERIAL CONTAMINATION AND METHOD OF USE

5 RELATED APPLICATION

This application claims priority to United States Provisional Application Serial No. 60/201,405, filed May 3, 2000.

FIELD OF THE INVENTION

10 The present invention relates to the detection of prokaryotic microorganisms. In particular, the present invention relates to a method and a device for detecting the presence or absence of a prokaryotic microorganism or pathogen that contaminates food and food related work areas.

15 BACKGROUND OF THE INVENTION

Presently, the United States Department of Food Safety and Inspection Services (FSIS) spend over half a billion dollars annually on meat, poultry, and fish inspections for bacterial contaminants. Broadly, the inspections are used to determine the cleanliness of the work area and to detect pathogenic microbes in foodstuffs. Ingestion of pathogenic
20 microbes can result in food poisoning when the microbes are inadvertently packaged into supermarket goods. One example of a pathogenic microbe is *Salmonella*. *Salmonella* is a genus of gram-negative bacterium that is a major source of human foodborne illness worldwide. Up to 4 million cases of salmonellosis are reported each year in the United States alone. A number of different serotypes of pathogenic *Salmonella* are known, the
25 most common of which are *S. typhimurium* and *S. enteritidis*. Typically, salmonellosis is treated with antibiotics. However, antibiotic resistant strains of *Salmonella* are emerging. *Listeria* and *E. coli* are also commonly occurring microbial contaminants. *Listeria monocytogenes* is a gram-positive bacterium that is a common cause of gastroenteritis. Tests for *E. coli* are performed as an indication of fecal contamination of food and work
30 areas.

The food industry has tended to test for food contamination at the production and wholesale level only. However, between the time a foodstuff is packaged and consumption occurs, a bacterial pathogen that was undetectable at the wholesale check can multiply and can thus become a health hazard. Contaminated food is increasing in importance for the producer. Bad publicity and product recalls cost the food industry millions of dollars each year.

Currently, the assay systems used at the wholesale and production level require specialized training and equipment, are time consuming, and are not very sensitive to the presence of small numbers of pathogenic bacteria. To assay for suspected bacterial contamination, a culture is obtained using a sample of the food or from the workplace that is suspected of being contaminated in order to increase the amount of detectable substance (suspected bacterial contaminant). Culturing may require up to 48 hours. Then, the cells grown in the culture are lysed to produce a lysate. Historically, assaying for pathogenic bacteria has been done using antibodies in an immunoassay for detecting bacterial proteins in the lysate. In recent years, some companies have used a polymerase chain reaction (PCR) based methods to replicate microbial genetic material for use in detection assays. The isolated genetic material is multiplied and identified in separate steps. The PCR detection system also requires hours rather than minutes to perform. Testing is performed at the time of or before shipping a product to the supermarket. A simple, sensitive, and rapid method for detecting food contamination is required both for testing at the wholesale level and at the retail level.

Detection devices are also needed at the retail level to protect the consumer. Testing apparatus useful at the wholesale level generally is not useful at the retail level for various reasons. First, most wholesale level tests require technical skills not often practiced by the typical consumer. Second, when a product remains on the shelf before purchase, contaminant levels can increase, especially if the temperature of the product is not appropriately controlled. During the time the food is in transit, bacterial growth can occur. Methods and devices for detecting microorganisms in food at the retail level have

been set forth. While tests have been developed for the retail market, they have failed to be adopted. Colorimetric detectors of changes in ionic content, for example pH, of the liquid or material surrounding the food product have been used on or incorporated into retail packaging such as illustrated in US Patent Nos. 4,746,616 and 5,053,339, the disclosures of each of which are incorporated herein by reference. Antibodies have also been coupled to colorimetric labels for detection of contaminants such as illustrated in US Patent Nos. 5,306,466 and 5,869,341, the disclosures of each of which are incorporated herein by reference. None of the aforementioned are sufficiently specific and sensitive to meet the needs of the current marketplace.

A means for specifically and sensitively detecting the presence or absence of at least one pathogenic microorganism in potentially contaminated food products at the retail level is needed.

SUMMARY OF THE INVENTION

The present invention provides a method for detecting the presence or absence of a prokaryotic microorganism comprising the steps of identifying a bacterial-specific protein, such as a protease that characterizes the presence of a specific prokaryotic microbe and thereby provides a marker for that microbe; detecting the protease that is a marker for the presence of a specific prokaryotic microbe by cleaving a substrate when the protease is present; and signaling the presence of that protease when cleavage has occurred. More specifically, the method comprises identifying at least one outer membrane protein or a secreted protein that is unique to a particular microbial pathogen such as for example *Listeria monocytogenes* and that is substrate specific.

Utilizing a bioinformatic approach, BLAST search comparisons of the protease complements of a plurality of common pathogenic bacterial species were made. One or more proteases that were unique to a species were identified. Next the specific substrate for each protease was identified using fluorescence resonance energy transfer (FRET).

The unique, identified substrate was then labeled and used to detect the presence of a

particular microbial pathogen in samples requiring testing. Identified protease substrates were tested with common foods and yeast in order to identify substrates that are specific to the bacteria.

5 In another aspect, the present invention provides a device for detecting the marker protein and for notifying a consumer of the presence of the marker protein. A biosensor for use in retail stores or home-use to detect contaminated food comprising a specific substrate that is coupled to packaging material proximal to the food stuff to be monitored for bacterial contamination is provided. Preferably, the substrate is covalently bound to a
10 label and thus has a detection signal that upon proteolysis of the substrate-label bond indicates the presence of the contaminating bacteria. The biosensor is made by first determining the specific substrate of a specific protease characteristic of the bacteria to be detected. The determined specific substrate is labeled with a plurality of chromatogenic or fluorescent leaving groups. Most preferably, the labeling group provides a latent
15 signal that is activated only when the signal is proteolytically detached from the packaging. Chromatogenic leaving groups are, for example, para-nitroanalide groups. Preferably, labeled substrate molecules are coupled to the interior surface of the packaging material proximal to the foodstuff. Should the foodstuff become contaminated, the microbial pathogen produces protease which acts on the labeled
20 substrate that is attached to the packaging, liberating the leaving group. When a visually detectable colorimetric signal is released, this results in a visible color change at the site of interest.

In another embodiment, proteases that are common among pathogenic microbial
25 species, but that are substrate specific and that share a common substrate, are identified for use in a method and a device to detect the presence of bacteria. For example, bacteria can collect on work surfaces in meat plants and in restaurants. A substrate that can be cleaved by a plurality of bacteria is labeled and covalently bound to a collector substrate, such as cotton fibers on the tip of a swab. The swab tip is used to wipe the surface
30 suspected in being contaminated by bacteria. The swab tip is placed in a medium and

incubated using conditions that allow proteolysis of the labeled substrate if the protease(s) specific for the bound, labeled substrate is present. Proteolysis results in the production of a detectable signal, for example a color change, may be visible if the signal is a chromogenic leaving group.

5

In yet another embodiment of the present invention, a strip sensor is provided. Preferably, the strip sensor comprises an inert material to which a substrate having a latent detection signal has been covalently attached. A single protease substrate or a plurality of protease substrates may be utilized. When a plurality of protease substrates are utilized, each may be labeled so as to distinguish it from another and/or each may be localized in a particular region on the strip support material. Depending upon the foodstuff, one can predict the frequency of expected occurrence of a particular pathogenic microbe. For example, *Listeria monocytogenes* is a common contaminant of poultry products, therefore, a chromagenic leaving group labeled substrate specific for a *Listeria monocytogenes* protease may be incorporated onto the surface of the packaging proximal to the poultry product. However, where multiple possible microbial pathogens may be present, a substrate that may be proteolyzed by all of the microbes may be utilized as long as it is specific to the pathogenic microbes.

10

15

20

In still another aspect of the present invention, a kit is provided for detecting microbial food contaminants. The kit comprises a flat solid support, preferably having a plurality of wells, to which a protein substrate labeled with leaving groups is linked. A means for providing a phosphate buffered solution, a negative control, and a positive control are provided. A sample of suspect food (SF) is prepared in phosphate buffer. An aliquot of each of SF, negative control, and positive control is placed in its own well and allowed to react. Those wells where a color change is observed contain microbial contaminant unless the negative control well also changes color. If the negative control well changes color, then the assay may be repeated.

25

30

BRIEF DESCRIPTION OF THE DRAWINGS

Figure 1 is an image of the Protein Encapsulated Styrofoam (PES) for capturing bacteria such as *L. monocytogenes*;

Figure 2 is a graph showing the dissolving time of PES tablets as determined by the amount of antibody released with time; and

Figure 3 is an image that demonstrates the fluorescence activation of a substrate in the presence of *L. monocytogenes*.

DETAILED DESCRIPTION

One of the most important considerations for developing a wholesale test kit is the ability to capture microbial pathogens in foods. It is necessary to have a procedure for reliably capturing bacteria from food matrices in such a way that all of the viable bacteria are captured and enriched in a suitable recovery medium. Many companies use cotton swabs or filters to capture the bacteria, but several independent studies have shown that entrapment of bacteria in these porous materials reduces the accuracy of detecting low-level contamination. Standard filter capture methods also entrap bacteria but are less reliable than straight-forward plating, and culture methods take up to a week to obtain results. Bacterial capture devices such as magnetic beads are more costly and require an additional step to collect the bacteria along the wall of the tube with a magnet.

In food safety testing, the food sample is ground up in a special blender called a stomacher blender, which is fitted with a sterile stomacher bag. The basic principle of the inventive capture method is to add a PES tablet to the blended sample for a few minutes with gentle shaking to capture the bacteria. The shell remains intact during this short exposure time at room temperature. After the bacteria are captured, the PES is allowed to float to the top of the stomacher bag from which it is removed with sterile forceps. The tablet is placed in growth enrichment medium and incubated at 37°C until sufficient bacteria can be reliably detected. After sufficient time at 37°C, the gelatin coat dissolves and the bacteria are effectively released into the growth culture medium.

To form the PES tablet, gelatin was heated to 65°C and allowed to cool to about 55°C . Then 50 µg/ml of antibody was added just before the gelatin capsule formed around the pellet. The PES technology can be used to capture and release bacteria from a test sample into enrichment medium. The rigidity of the gelatin prevents the antibodies from being released until several hours after incubation (*see* figure 2).

Two PES tablets were incubated in PBS with vigorous shaking at room temperature (RT; series 1 and 2). At each time point (0, 5, 10, 20, 60, 120, and 720 minutes), an aliquot of the sample was removed and the amount of HRP conjugated antibody was detected with TMB-peroxidase substrate (KPL, Gaithersburg, MD). The samples were read with a Benchmark microplate reader at 655 nm. After 20 minutes, less than 1% of the total antibody on the PES tablet has dissolved into the PBS solution. Thus it is possible to capture bacteria with the antibody-coated capsules and release them into the enrichment media with this method. The rigidity (high bloom number) of the pig skin gelatin (300 bloom) makes this capture and release procedure possible. Using a lower bloom gelatin would allow for the bacteria to be released faster into the enrichment media.

The present invention provides a method to quantitatively capture and release contaminating bacteria present in a food sample that utilizes the protein-encapsulated Styrofoam (PES) tablets described above. One or more tablets coated with commercially available antibodies entrapped in gelatin can be placed in a homogenized food test sample in a stomacher bag and incubated with gentle shaking at room temperature. The tablet can then be removed with sterile forceps and placed in growth medium at 37°C for a short period of time, until the gelatin coat dissolved and released the bacteria into the medium.

The presence of *L. monocytogenes* contamination is detected by measuring the metalloprotease (mpl), which is found only in the pathogenic species of *L. monocytogenes* (Domann et al., Infection and Immunity. 1991. Jan;59(1):65-72; Mengaud et al., 1991). A series of peptides, M1 (lot 038-13/15) and P1 (lot 038-10/12),

for use as synthetic substrates for mpl (Table 1) was designed and the sequences manufactured by New England Peptide (195 Kimball St., Fitchburg, MA). The peptides were synthesized with fluorescent groups (DABCYL and EDANS) in such a way that the active sites were not blocked. The M1 peptide is specific for *L. monocytogenes* at pH 5.5.

5

Table 1: First set of peptides used as probes for FRET analysis of mpl activity

Peptide	Amino acid sequence	MW	Description
M1	DABSYL-NMLSEVERE-EDANS	1642	Specific to <i>Listeria</i> at pH 5.5
P1	DABSYL-ACCDEYLQTKE-EDANS	1838	Substrate with broad pathogen specificity
P2	DABSYL-ADTVEPTGAKE-EDANS	1653	Not cleaved by <i>Listeria</i> or <i>E. coli</i>

(peptide M1 = SEQ ID NO. 1; P1 = SEQ ID NO. 2; P2 = SEQ ID NO. 3)

Preliminary evidence indicates that at pH 5.5, the M1 peptide is specific for *L. monocytogenes* and does not significantly detect the presence of *E. coli* (see Figure 3). It is expected that this peptide and derivatives of this peptide (with one or two amino acid substitutions) could be used for a standard microplate assay format for *L. monocytogenes* in foods.

It was discovered that *Listeria monocytogenes* secretes at least two characteristic proteases, one of which is a metalloprotease and one that is a serine protease.

Microbial proteases can be separated into four major classes based on the catalytic site (metallo, serine, cysteine, and aspartate proteases). The particular type of catalytic site may be determined using catalytic site blockers. For example the presence of a metalloprotease having a metallo catalytic site can be demonstrated by specifically blocking the catalyzed reaction with TAPI [N-{D,L-[2-(hydroxaminocarbonyl)m-ethyl]-4-methylpentanoyl}L-3-(2(naphthyl)alanyl-L-alanine, 2-aminoethyl amide]. The catalytic site of each protease has a characteristic amino acid signature, motif, or amino acid sequence that can be used to identify that protease and similar new proteases within a class thereof. In addition, many of the proteases have two forms: a larger inactive form that requires partial proteolysis (limited proteolysis) to obtain a smaller active form of the

enzyme. Proteases are known to break the peptide bonds in a protein substrate. Some proteases are known to be very specific with regard to the type of peptide bond that they are able to hydrolyze or break. When a protease is limited with respect to the bond(s) it can break, the protease is said to be substrate specific.

5

The amino acid sequence characteristic of a particular enzyme or of its catalytic site is reproduced from the genetic code. Knowing the amino acid sequence of an enzyme or of its catalytic site, one can deduce the nucleotide sequence in the DNA or RNA that codes for that sequence. The nucleotide sequence can then be located in the genetic code of a bacterium or it can be synthesized. Cloning of the genetic sequence coding for the amino acid sequence can be used to provide multiple templates for generating copies of the enzyme or its catalytic site. Using these copies of the enzyme or its catalytic site, the exact site of cleavage or type of bond hydrolyzed can be determined. Using this information, a specific substrate of the enzyme can be defined. This substrate can be used to assay for the presence of the pathogenic bacteria. The substrate is provided with a label that serves to signal the presence of the pathogenic bacteria. The preferred label is chromagenic and produces a visual signal when cleaved. For example, when determining specific cleavage of a substrate, the substrate can be provided with two different dyes, where one serves to quench the other when the dyes are in close proximity and measured by FRET.

10

15

20

Fluorescence resonance energy transfer (FRET) is the process of a distance dependent excited state interaction in which the emission of one fluorescent molecule is coupled to the excitation of another. A typical acceptor and donor pair for resonance energy transfer consists of 4-[[4-(dimethylamino) phenyl]azo] benzoic acid (DABCYL) and 5-[(2-aminoethylamino)naphthalene sulfonic acid (EDANS). EDANS is excited by illumination with 336 nm light, and emits a photon with wavelength 490 nm. If a DABCYL moiety is located within 20 angstroms of the EDANS, this photon will be efficiently absorbed. DABCYL and EDANS will be attached to opposite ends of a peptide substrate. If the peptide is intact, FRET will be very efficient. If the peptide has

25

30

been cleaved by the metalloprotease, the two dyes will no longer be in close proximity and FRET will be inefficient. The cleavage reaction can be followed by observing either a decrease in DABCYL fluorescence or an increase in EDANS fluorescence (loss of quenching).

5

Alternative methods for providing a visual signal include labeling the protease substrate with a food safe dye (FD&C) such as Texas red or with amido black or with a fluorometric dye such as 4-nitroaniline or 7-amino-4-methyl coumarin. For retail applications, labeled substrate may be bound to the food packaging. When the protease characterizing the pathogenic microorganism is present, the label or the labeled substrate is cleaved from the packaging, diffusing from its attached site. This results in a color change at that site. For example, the packaging may have "DO NOT EAT" printed in red thereon. When the labeled substrate is present, the writing is masked. When the label is removed, the warning becomes visible.

10

15

In one embodiment, an assay system that detects and signals the presence of pre-determined prokaryotic microorganisms is provided. Detection of *Listeria monocytogenes* is exemplified. However, the invention is not intended to be limited to that pathogenic species. The presence of the pre-determined prokaryotic organism of interest is indicated by proteolytic cleavage of a substrate by a protease that results in the production of a signal. Preferably, the signal is chromogenic. *L. monocytogenes* (DP-L1955) was kindly provided by Dr. Julie Theriot of the Stanford School of Medicine. The *L. monocytogenes* was made avirulent by a deletion of the gene required for intracellular motility (ActA, Smith et al., J Cell Biol. 1996 Nov;135(3):647-60).

20

25

A method for developing an assay for detecting a pathogenic bacteria that produces at least one extracellular protease and a method for using the assay to detect pathogenic bacteria producing at least one extracellular protease follows:

Step 1) Define a catalytic marker amino acid sequence that uniquely identifies the prokaryotic microorganism of interest.

Select an amino acid sequence, also termed a marker sequence, that uniquely characterizes or marks the presence of the microorganism of interest, such as, for example, an amino acid sequence unique to an extracellular protease produced by a pathogenic prokaryotic microorganism of interest. The selection is performed utilizing a bioinformatic approach. BLAST search comparisons among common pathogenic bacterial species with respect to known amino acid sequences are made. One or more amino acid sequences that are unique to a specific prokaryotic microorganism are determined.

Step 2) Obtain sufficient protease to determine conditions facilitating optimal proteolysis by the protease.

Isolate the protease from the extracellular medium in which the pathogenic bacteria to be assayed is growing.

Alternatively, if the genetic sequence for the protease or the location of the genetic sequence for the protease are unknown, isolate and clone the genetic sequence that results in the expression of the marker amino acid sequence of Step 1, or, first determine the genetic sequence, and then proceed as before.

Step 3) Determine the conditions for growth of the prokaryotic organism and for the production of extracellular protease.

Determine medium required for growth of the specific prokaryotic microorganism of interest and for expression of its unique active protease into the medium.

Determine whether a second protease is required to convert the specific protease from an inactive precursor form to an active form.

To determine if the protease has been secreted in an active form, a sample of the bacterial culture is provided with chosen potential substrates and cleavage of these substrates is determined. For example, a bacteria pellet from Step 3 is gently resuspended in 500 µl of DMEM in the presence of common extended substrates (myosin II, bovine serum albumin, collagen, fibronectin, and hemoglobin). Myosin II, but not the proteins from milk such as casein, has been shown to be a suitable substrate to measure the proteolytic activity of gram positive bacteria isolated from dried cured ham (Rodriguez et al., J Appl Microbiol. 1998. Nov;85(5):905-12). The suspension of bacteria and substrates is incubated at 37°C with gentle shaking. At preset times (0.1, 0.3, 1.0, 3.0, 5.0, 24, and 48 hours) the samples are centrifuged to spin down the bacteria, and a small aliquot is removed for a SDS-PAGE gel sample. After completion of the time course the samples are run on a 10-15% gradient SDS-PAGE minigel. Then, the proteins are transferred to Immobilon Pseq (Transfer buffer, 10% CAPS, 10% methanol pH 11.0, 15 V for 30 minutes) using a Bio-Rad semi-dry transblotting apparatus. Following transfer of the proteins, the blot is stained with Coomassie blue R-250 (0.25% Coomassie Brilliant Blue R-250, 50% methanol, 10% acetic acid) and destained (high destain for 5 minutes, 50% methanol, 10% acetic acid; low destain until complete, 10% methanol, 10% acetic acid) followed by sequencing from the N-terminal. Alternatively, the samples will be run on a mass spectrometer in order to map the sites of proteolytic cleavage using a Voyager Elite Mass spectrometer (Perceptive Biosystems).

For example, an extracellular metalloprotease is produced by *L. monocytogenes*, designated mpl. The gene for this protease is just downstream of another virulence factor, the listerolysin gene (Domann et al., Infection and Immunity. 1991. Jan;59(1):65-72). Mpl has two forms, active and inactive. The three possible mechanisms for activation of mpl or any inactive protease by limited proteolysis are: auto-digestion (autocatalytic processing), digestion by a bacterial protease other than the

metalloprotease, and digestion by an extracellular host cell protease. Metalloprotease activity has been found to be indirectly measurable in the supernatants of *L.*

monocytogenes. Mpl activity can be monitored by measuring the shedding of IL-6 receptor from the surface of human monocytes. Processing of the precursor

5 metalloprotease of *L. monocytogenes* to the mature form was determined to require the presence of the monocytes.

The following method can be used to test activation of an inactive protease of other pathogenic bacteria. Preincubate the bacteria of interest with either methanol fixed
10 whole cells or NaOH stripped membrane preparations of monocytes. Fixation of the monocytes will prevent the bacteria from internalizing. Briefly, 10,000 CFU of *L. monocytogenes* is added to nearly confluent dishes of monocytes that are fixed with minus 20°C 100% methanol for 10 minutes or 5.0% paraformaldehyde in a buffer that preserves the cytoskeleton. After 2-4 hours of incubation, the supernatant containing
15 bacterium is removed and is centrifuged to separate the supernatant from the cell pellet. The supernatant is incubated with chosen protein substrates for preset times to determine its activity.

L. monocytogenes expression of active metalloprotease was found to be enhanced
20 by growing the prokaryotic bacteria in a protein free medium such as D10 medium, a medium developed by Trivett and Meyer (1971). The bacteria were cultured overnight (24 hrs); centrifuged at 5K x g for 10 minutes and then resuspended in the tissue culture medium (Dulbecco's Modified Eagle Medium, "DMEM", Life Technologies (Gaithersburg, MD).

25
Step 4) Identify any specific protein and/or peptide substrate(s) of the active species-specific protease.

Step 5) Increase the specificity of the protease-substrate interaction (optional) by
30 determining the active or binding site of the protease, then determining the genetic

sequence useful for producing that active or binding site, and cloning the determined genetic sequence.

Identify the target site(s) also termed the "active site(s)" for cleavage by the protease.

Determine the peptide sequence for the active site of the protease using fluorescence resonance energy transfer (FRET). Fluorescence resonance energy transfer (FRET) is the process of a distance dependent excited state interaction in which the emission of one fluorescent molecule is coupled to the excitation of another. FRET is one of the best ways to identify the substrate specificity for a particular protease (Liu et al., Anal Biochem. 1999. Feb 15;267(2):331-5). A typical acceptor and donor pair for resonance energy transfer consists of 4-[[4'-(dimethylamino) phenyl]azo] benzoic acid (DABCYL) and 5-[(2-aminooethylamino)-naphthalene sulfonic acid (EDANS). EDANS is excited by illumination with 336 nm light, and emits a photon with wavelength 490 nm. If a DABCYL moiety is located within 20 angstroms of the EDANS, this photon will be efficiently absorbed. DABCYL and EDANS will be attached to opposite ends of a peptide substrate. If the peptide is intact, FRET will be very efficient. If the peptide has been cleaved by the metalloprotease, the two dyes will no longer be in close proximity and FRET will be inefficient. The cleavage reaction can be followed by observing either a decrease in DABCYL fluorescence or an increase in EDANS fluorescence (loss of quenching).

When the enzyme is a metalloprotease (mpl), the active form of the mpl can be amplified by polymerase chain reaction using forward and reverse primers containing useful restriction sites for cloning into an expression vector with a strong inducible promoter (such as the T7 system of pET-28a, Novagen). Useful primers for the amplification are catgccatgggtagaacgggctgataccca (SEQ ID NO. 4) and gcgcccgaattctcagttaaccccaactgctt (SEQ ID NO. 5). Each primer has at least a 6 base pair overhang. The DNA is amplified from either lysed colonies of *L. monocytogenes* or from

purified preparations of genomic DNA from *L. monocytogenes*. The forward primer is provided with a *Nco* I site that includes the start methionine (CATATG). The reverse primer includes a stop codon (TGA) and an *Eco* RI restriction site (GAATTC). A typical amplification protocol comprises amplifying the genetic sequence with Taq polymerase under the following conditions: 92°C initial melt 4 minutes (25 cycles, 50°C anneal 30 seconds, 72°C extension 1.5 minutes, 92°C melt 30 seconds), 72°C final extension 10 minutes and 4°C store. The annealing temperature may be increased or decreased somewhat to account for the predicted melting temperature (T_m) of the primers. Following cloning of the genetic sequence for the active metalloprotease into pET28a, the gene is sequenced by double stranded primer walk sequencing on an ABI310 genetic analyzer. The recombinant active form of the metalloprotease genetic sequence is purified by nickel-nitrilotriacetic acid chromatography by taking advantage of the polyhistidine tag in the pET-28a vector.

Step 6) Provide a device with a peptide substrate having a labeled carboxy terminus where cleavage thereof has been determined to be specific by the FRET assay of Step 5 to the protease of the pathogenic bacteria of interest.

For example, a chromagenically labeled peptide could be bound to a PVDF membrane with a hydrophobic area. When the peptide is clipped, the dye would concentrate in the hydrophobic area, becoming visible. Alternatively, the peptide could be labeled fluorescently, and visualized with a UV lamp. Another possibility is a peptide with a chromagenic leaving group that upon removal changes color. The peptide may be labeled with a blue dye, coupled to a red surface. To the observer, the surface would look black unless the specific bacteria were present to provide protease that would cleave the peptide, turning the surface red as the peptide was proteolysed and diffused away. Although conventional leaving groups such as 4-nitroanaline or 7-amino-4-methylcoumarin can be quantified spectrophotometrically and fluorometrically, it may be best to use a leaving group such as Texas Red that has an intense blue color. By placing two or

three Texas Red groups near the end of the peptide, it would be possible to detect visually the clearance of the substrate by *L. monocytogenes* protease activity.

The cysteines at carboxyl terminus of the peptides were labeled with Texas Red bromoacetamide using a standard procedure described for labeling actin. See Wang and Sanders, 1990. An example is a hydrophobic peptide labeled with 2-3 Texas Red molecules. Another possibility would be to label the peptide with Amido black. As the substrate is proteolysed by mpl, the black peptide diffuses away, and a clear zone would appear indicating the presence of *L. monocytogenes*.

The peptide can be coupled to a surface such as a glass microtiter plate or a polypropylene food packaging material in the proper orientation. The microtiter plate should provide a plurality of derivatized binding sites for coupling to the peptide substrate such as succinidyl ester labeled primary amine sites on derivatized plates (Xenobind plates, Xenopore).

By maintaining a neutral pH, the amino terminus of the peptide can be preferentially labeled. Briefly, the peptide (at a concentration of 1-3 mg/ml) will be incubated with a Xenobind plates for 2 hours in 50 mM sodium phosphate buffer, (pH 7.0). Following rinsing with wash buffer, the plates will be blocked with 3% bovine serum albumin for 2 hours at 37°C. Finally the plates will be rinsed three times in wash buffer and stored at 4°C until use.

Optionally unoccupied reactive sites on the microtiter plate or on the food packaging are blocked by coupling bovine serum albumin, or the active domain of p26 thereto. p26 is an alpha-crystallin type protein that is used in this case to reduce non specific protein aggregation. The ability of the p26 protein to refold heat denatured citrate synthetase before and after coupling to the surface of the food packaging is used as a control for determining p26 activity. Alpha-crystallin type proteins were recombinantly produced using standard recombinant DNA technologies (Maniatis et al., Molecular

cloning: a laboratory manual. 1982). Briefly, the plasmid containing the beta sheet-charged core domain of p26 is electroporated into electrocompetent B121(DE3) cells (Bio-Rad E. coli pulser). The cells are grown up to an OD of 0.8, then induced with 1mM IPTG for 4 hours. The cells are spun down, and sonicated in low buffer (10 mM Tris, pH 8.0, 500 mM NaCl, 50mM Imidazole) to lyse (Virsonic, Virtis, Gardiner, NY). The lysate is spun down at 13,000 x g for 10 minutes, and the supernatant 0.45 and 0.2 µm filtered. This filtrate is loaded onto a Ni-NTA superose column (Qiagen, Valencia, CA, cat # 30410). High buffer (10 mM Tris pH 8.0, 500 mM NaCl, 250 mM Imidazole) is used to elute the protein.

10

EXAMPLE 1

Rapid *Listeria monocytogenes* Detection at the Wholesale Level

Food products were tested in pentuplicate to determine the range of applicability for ready to eat food products. Each solid food sample was processed in a laminar flow hood. Twenty-five grams of each food product were removed from its package and weighed. The weighed sample of the solid food was placed in a sterile stomacher bag with 125 ml of phosphate-buffered saline (PBS, 10 mM potassium phosphate, pH 7.4, 120 mM sodium chloride, 2.7 mM potassium chloride). Solid food was homogenized in a stomacher blender (Fisher Scientific; Pittsburgh, PA) and the filtered supernatants were used for testing. *L. monocytogenes* was captured and efficiently released by adding a plurality of Protein Encapsulated Styrofoam Tablets. PES tablets (described above) were incubated with the food extract for at least a 30 minute incubation period. The PES tablet were removed from the surface and placed in enrichment media to efficiently release the bacteria. After incubating at 37°C for at least 4 hours, the media was spun down at 5,000xg for 10 minutes, and the pellet was resuspended in 50 µl of PBS. Liquid foods were prepared by mixing an aliquot of the liquid food with one of the PES tablets without stomaching the liquid. After the 30 minute capture step, the PES tablet was recovered and the pellet was placed in enrichment media.

30

A protease based assay was performed on each sample. Each well in the microtiter plate was read before starting the assay in order to get an accurate measurement of the background to control for well to well variability. The protease assay was performed using microtiter plates having FRET labeled peptide such as M1 that is described above. Typically 0.5 µg of peptide was incubated with 50µl of bacterial culture extract for at least 20 min at 37°C. In the presence of bacteria the FRET peptide will fluoresce at 493 nm. As positive controls, either *L. monocytogenes* or the active recombinant form of the pathogen specific protease can be added to some of the food samples prior to incubation in the microtiter plate assay. Negative controls consist of food samples that are determined to be devoid of bacterial pathogens based on existing published methods (Curtis et al., 1995).

EXAMPLE 2

Rapid *Listeria monocytogenes* Detection of Bacteria for Home Use

PVDF is a blotting membrane that is hydrophobic until soaked in methanol, when it becomes hydrophilic. A ring of methanol was stamped on PVDF and allowed to air dry. A peptide substrate labeled with an FD&C approved dye was incubated with the membrane. It only bound to the hydrophilic ring. This membrane was attached to packaging material, for example the inside of food packages in direct contact with the food. If the peptide is cleaved by bacterial proteases in the food, the dye molecule will be attracted to the hydrophobic region of the membrane, and concentrate there, indicating that bacteria is present by the formation of color or florescence within the center of the ring. Bacteria other than *Listeria monocytogenes* can be detected using the described method and device once a specific protease substrate interaction has been determined and optimized for the bacterium of interest.

For some bacterial species, it may be necessary to provide a method of optimizing substrate-protease interaction. For example, divalent cations may be required to optimize protease cleavage of the coupled substrate. In this case, the food packaging surface will

be blocked with proteins and/or peptides that can be precharged with metal ions, such as for example polyhistidine or zinc finger motifs.

One of ordinary skill in the art will appreciate further features and advantage of
5 the invention based on the above-described embodiments. Accordingly, the invention is not to be limited by what has been particularly shown and described, except as indicated by the appended claims. All publications and references cited herein are expressly incorporated herein by reference in their entirety

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
2201
2202
2203
2204
2205
2206
2207
2208
2209
2210
2211
2212
2213
2214
2215
2216
2217
2218
2219
2220
2221
2222
2